## **CASE SUMMARY**

Case #2011/1081

**Complainant:** Bart's Water Sports

7581 East 800 North

North Webster, Indiana 46555

574-834-7666


**Applicator:** Chad Tom

Tippy View Lawn Care 6590 North Kalorama Road Leesburg, Indiana 46538

574-527-9373

Licensed Applicator Licensed Business

- 1. On June 28, 2011, I, Agent Kevin Gibson of the Office of Indiana State Chemist (OISC), performed an investigation at the complainant's property in response to a claim of injury/damage to non-target trees possibly resulting from exposure to the herbicide Imprelis. A Notice of Inspection was issued to Bart Culver. I observed the following during my on-site investigation:
  - a) Spruce tree with brown needles throughout tree and close-up (see figures #1 & #2).
  - b) Spruce tree browning needles and close-up (see figures #3 & #4).
  - c) Ginko tree and close-up (see figures #5 & #6).
- 2. I took the following photos depicting injured/damaged vegetation:



- 3. I collected the following vegetation samples from visibly impacted non-target vegetation as described in paragraph #1 for examination by the Purdue Plant Pest Diagnostic Laboratory (PPDL):
  - a) Spruce
  - b) Ginko

, Hebron

- 4. I collected the following environmental samples for chemical analysis by the OISC Residue Laboratory:
  - a) Composite vegetation sample from yard (spruce)
  - b) Vegetation sample from yard (ginko)
  - c) Composite soil sample from yard (inside drip line)
- 5. According to a report from the PPDL, "There was no evidence of significant disease or mite/insect injury on the samples submitted. The sample (and pictures) submitted show symptoms that are typically found to be associated with injury caused by synthetic auxin (growth regulator type) herbicides. Typical symptoms caused by these herbicides can include epinasty (twisting and curving) of the leaves or needles, shoot and shoot tip; leaf cupping which can be upward or downward, and in extreme cases, new leaves can be irregular in size and shape (usually smaller than normal) and have abnormal leaf margins."
- 6. According to the report from the OISC Residue Lab the following levels of aminocyclopyrachlor (active ingredient in Imprelis Herbicide) were found in the samples referenced in item #4:

a) Composite vegetation sample (spruce)
 b) Vegetation sample (ginko)
 c) Composite soil sample (outside drip line)
 PPB=Parts Per Billion

7. According to the application information collected from the applicator Imprelis Herbicide (EPA Reg. No. 352-793) was applied on May 3, 2011, at the rate of 4.3 oz. per acre using ground spray equipment; no application was made to the soil within the drip line of any of the trees or ornamentals; no application was made directly to any exposed roots of any trees or ornamentals.

Kevin W. Gibson Date: October 3, 2011
Pesticide Investigator

**Disposition:** No violation of the Indiana Pesticide Use and Application Law was documented against the pesticide applicator. Effective September 15, 2011, the Indiana registration for Imprelis Herbicide, EPA Reg. #352-793, was cancelled because it was determined by OISC that the product is "misbranded" (it bears label directions that are inadequate to prevent unreasonable adverse effects to non-target vegetation).

George N. Saxton Final Date: October 20, 2011

Compliance Officer